AD-A065 849 FOREIGN TECHNOLOGY DIV WRIGHT-PATTERSON AFB OHIO F/G 20/4
UNSTEADY FLOW OF A VISCOUS FLUID,(U)
JUL 78 A N GOTS, G K BERMAN, K V VALIKOV
UNCLASSIFIED FTD-ID(RS)T-1110-78 NL

| OF |

AD
A065849

END
DATE
FILMED
5-79
DDC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A065849

FTD-ID(RS)T-1110-78

1

FOREIGN TECHNOLOGY DIVISION

UNSTEADY FLOW OF A VISCOUS FLUID

By

A. N. Gots, G. K. Berman and K. V. Valikov

DDC
RECEIVED
MAR 19 1979
A

Approved for public release;
distribution unlimited.

78 12 22 235

EDITED TRANSLATION

FTD-ID(RS)T-1110-78 17 July 1978

MICROFICHE NR: FTD-78-C-000985

CSB77011787

UNSTEADY FLOW OF A VISCOUS FLUID

By: A. N. Gots, G. K. Berman and K. V. Valikov

English pages: 3

Source: Sbornik Nauchnykh Trudov Ivanovskiy Energeticheskiy Institut Im. V. I. Lenina, Issue 20, Ivanovo-Vladimir, 1972, pp. 162-163

Country of Origin: USSR
Translated by: Carol S. Nack
Requester: FTD/TQTA
Approved for public release; distribution unlimited.

A

THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WP-AFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES TRANSLITERATION SYSTEM

Block	Italic	Transliteration	Block	Italic	Transliteration
А а	*А а*	A, a	Р р	*Р р*	R, r
Б б	*Б б*	B, b	С с	*С с*	S, s
В в	*В в*	V, v	Т т	*Т т*	T, t
Г г	*Г г*	G, g	У у	*У у*	U, u
Д д	*Д д*	D, d	Ф ф	*Ф ф*	F, f
Е е	*Е е*	Ye, ye; E, e*	Х х	*Х х*	Kh, kh
Ж ж	*Ж ж*	Zh, zh	Ц ц	*Ц ц*	Ts, ts
З з	*З з*	Z, z	Ч ч	*Ч ч*	Ch, ch
И и	*И и*	I, i	Ш ш	*Ш ш*	Sh, sh
Й й	*Й й*	Y, y	Щ щ	*Щ щ*	Shch, shch
К к	*К к*	K, k	Ъ ъ	*Ъ ъ*	"
Л л	*Л л*	L, l	Ы ы	*Ы ы*	Y, y
М м	*М м*	M, m	Ь ь	*Ь ь*	'
Н н	*Н н*	N, n	Э э	*Э э*	E, e
О о	*О о*	O, o	Ю ю	*Ю ю*	Yu, yu
П п	*П п*	P, p	Я я	*Я я*	Ya, ya

*ye initially, after vowels, and after ъ, ь; e elsewhere.
When written as ё in Russian, transliterate as yё or ё.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	$\sinh^{-1}$
cos	cos	ch	cosh	arc ch	$\cosh^{-1}$
tg	tan	th	tanh	arc th	$\tanh^{-1}$
ctg	cot	cth	coth	arc cth	$\coth^{-1}$
sec	sec	sch	sech	arc sch	sech^{-1}
cosec	csc	csch	csch	arc csch	csch^{-1}

Russian	English
rot	curl
lg	log

1110

UNSTEADY FLOW OF A VISCOUS FLUID

A. N. Gots, G. K. Berman and K. V. Valikov

We will consider the unsteady flow of a viscous fluid in a rectangular channel. We will direct the z-axis along the channel, whose width will be designated as a, and height - b. We will consider the flow conditions to be isothermic and unsteady. In this case, the differential equation of motion [1] is

$$\frac{\partial v}{\partial t}=\frac{f(t,z)}{\rho}+\nu\left(\frac{\partial^2 v}{\partial x^2}+\frac{\partial^2 v}{\partial y^2}\right). \qquad (1)$$

where v is the flow rate on the z-axis; ρ is the density of the fluid; $f(t, z)=-\frac{\partial P}{\partial z}$ is the pressure gradient; $\nu=\frac{\eta}{\rho}$ is the kinematic viscosity coefficient; and η is the viscosity of the fluid.

Equation (1) was written with the assumption that $v_x = v_y = 0$; v_z = v(x, y, z). The initial and boundary conditions are

$$v|_{t=0} = 0;\ v|_{\Gamma} = 0, \qquad (2)$$

where Γ is the cross-sectional profile.

We will search for the solution to differential equation (1) with conditions (2) in the form [2]

$$v(x, y, t) = \sum_{n,m=0}^{\infty} v_{nm}(t) \sin \frac{\pi(2n+1)x}{a} \sin \frac{\pi(2m+1)y}{b}, \qquad (3)$$

where v_{nm} is the coefficient to be determined.

We will expand function f(t, z) into a binary Fourier series, after the calculation of the coefficients of which we will have

$$f(t, z) = \frac{16}{\pi^2} \sum_{n,m=0}^{\infty} \frac{1}{(2n+1)(2m+1)} \sin \frac{\pi(2n+1)x}{a} \sin \frac{\pi(2m+1)y}{b}. \qquad (4)$$

Substituting expressions (3) and (4) in equation (1), and also considering conditions (2), we will find

$$v_{nm}(t) = \frac{16}{\pi^2 \rho (2n+1)(2m+1)} \int_0^t e^{-\nu\left[\frac{(2n+1)^2}{a^2} + \frac{(2m+1)^2}{b^2}\right](t-\tau)} f(z, \tau)\, d\tau. \qquad (5)$$

Thus, solutions (3) and (5) obtained make it possible to find

the law of the distribution of the velocity of a viscous fluid in a rectangular channel.

Bibliography

1. N. A. Slezkin. Dynamics of a Viscous Incompressible Fluid, Gostekhizdat, M., 1965.

2. N. S. Koshlyakov, E. B. Gliner, M. M. Smirnov. Main Differential Equations of Mathematical Physics, Fizmatgiz, M., 1962.

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

ORGANIZATION	MICROFICHE
A205 DMATC	1
A210 DMAAC	2
B344 DIA/RDS-3C	9
C043 USAMIIA	1
C509 BALLISTIC RES LABS	1
C510 AIR MOBILITY R&D LAB/FIO	1
C513 PICATINNY ARSENAL	1
C535 AVIATION SYS COMD	1
C591 FSTC	5
C619 MIA REDSTONE	1
D008 NISC	1
H300 USAICE (USAREUR)	1
P005 ERDA	1
P005 CIA/CRS/ADD/SD	1
NAVORDSTA (50L)	1
NASA/KSI	1
AFIT/LD	1

ORGANIZATION	MICROFICHE
E053 AF/INAKA	1
E017 AF/RDXTR-W	1
E403 AFSC/INA	1
E404 AEDC	1
E408 AFWL	1
E410 ADTC	1
E413 ESD	2
FTD	
CCN	1
ASD/FTD/NICD	3
NIA/PHS	1
NICD	2